

Cardiac anti-remodelling effect of aerobic training is associated with a reduction in calcineurin/NFAT signalling pathway in heart failure mice

R. S. F. Oliveira¹, J. C. B. Ferreira¹, E. R. M. Gomes², N. A. Paixão¹, N. P. L. Rolim¹, A. Medeiros¹, S. Guatimosim², P. C. Brum¹

¹School of Physical Education and Sport, University of Sao Paulo, SP, Brazil and

²Department of Physiology and Biophysics, Federal University of Minas Gerais, MG, Brazil.

Running head: Exercise training attenuates calcineurin signalling in heart failure

Author for correspondence:

Patricia Chakur Brum, PhD.

Escola de Educação Física e Esporte da Universidade de São Paulo

Departamento de Biodinâmica do Movimento do Corpo Humano

Av. Professor Mello Moraes, 65 - Butantã - São Paulo – SP

CEP 05508-900– BRASIL

Telephone: (5511) 3091-3136 – FAX: (5511) 3813-5921

e-mail: pcbrum@usp.br

Abstract

Cardiomyocyte hypertrophy occurs in response to a variety of physiological and pathological stimuli. While pathological hypertrophy in heart failure is usually coupled with depressed contractile function, physiological hypertrophy associates with increased contractility. In the present study, we explored whether 8-weeks of moderate intensity exercise training would lead to a cardiac anti-remodelling effect in an experimental model of heart failure associated with a deactivation of a pathological (calcineurin/NFAT, CaMKII/HDAC) or activation of a physiological (Akt-mTOR) hypertrophy signalling pathway. The cardiac dysfunction, exercise intolerance, left ventricle dilation, increased heart weight and cardiomyocyte hypertrophy from mice lacking α_{2A} and α_{2C} adrenoceptors (α_{2A}/α_{2C} ARKO mice) were associated with sympathetic hyperactivity induced-heart failure. The relative contribution of Ca^{2+} -calmodulin high-affinity (calcineurin/NFAT) and low-affinity (CaMKII/HDAC) targets to pathological hypertrophy of α_{2A}/α_{2C} ARKO mice was verified. While nuclear calcineurin B, NFATc3 and GATA-4 translocation were significantly increased in α_{2A}/α_{2C} ARKO mice, no changes was observed in CaMKII/HDAC activation. As expected, cyclosporine treatment decreased nuclear translocation of calcineurin/NFAT in α_{2A}/α_{2C} ARKO mice, which was associated with improved ventricular function and pronounced anti-remodelling effect. Akt-mTOR signalling pathway was not activated in α_{2A}/α_{2C} ARKO mice. Exercise training improved cardiac function and exercise capacity in α_{2A}/α_{2C} ARKO mice and decreased heart weight and cardiomyocyte width paralleled by diminished nuclear NFATc3 and GATA-4 translocation as well as GATA-4 expression levels. Combined, these findings support the notion that deactivation of calcineurin/NFAT pathway-induced pathological hypertrophy is a preferential mechanism by which exercise training leads to cardiac anti-remodelling effect in heart failure.

Key words: aerobic exercise training, cardiac hypertrophy, heart failure, calcineurin-NFAT pathway, Akt-mTOR pathway.

Abbreviations list

Akt, protein kinase B; α_{2A}/α_{2C} ARKO, α_{2A} and α_{2C} adrenergic receptor knockout mice; β -MHC, β -myosin heavy chain; CaMKII, Ca^{2+} /calmodulin-dependent protein kinase II; FS, fractional shortening; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; GATA-4, GATA binding protein 4; HDAC 4 and 5, class II histone deacetylases; IGF-I, insulin-like growth factor; LVEDD, left ventricular end-diastolic dimension; LVESD, left ventricular end-systolic dimension; mTOR, mammalian target of rapamycin; NFATc3, nuclear factor of activated T-cells transcription factor; PI3K, phosphatidylinositol 3-kinase; SERCA2, sarcoplasmic reticulum Ca^{2+} -ATPase; WT, wild type.

Introduction

Cardiac hypertrophy is an adaptive response of the heart to a variety of pathophysiological stimuli, such as hypertension, myocardial infarction, valvular insufficiency, infectious agents, or mutations of contractile proteins. Pathological hypertrophy is associated with severe cardiac dysfunction, arrhythmias, sudden death, and heart failure (Levy *et al.*, 1990; Frey & Olson, 2003). Indeed, epidemiological studies have revealed that cardiac hypertrophy is an independent risk factor for heart failure development (Ho *et al.*, 1993; Lorell & Carabello, 2000). However, not all forms of cardiac hypertrophy are pathological, since exercise training induces physiological cardiac hypertrophy associated with improved cardiac function in athletes (Naylor *et al.*, 2008).

Several studies have reported Ca^{2+} -handling abnormalities in hypertrophied and failing myocardium in response to neurohumoral activation, stretch and pacing (Bustamante *et al.*, 1991; Balke & Shorofsky, 1998; Rossman *et al.*, 2004; MacDonnell *et al.*, 2007; Rolim *et al.*, 2007). In fact, decreased Ca^{2+} transient peak and prolonged Ca^{2+} decay with increased diastolic intracellular Ca^{2+} have been described in animal and human heart failure (Gwathmey *et al.*, 1987; Schwinger *et al.*, 1995; Seki *et al.*, 2003; Bartholomeu *et al.*, 2008), and related to sustained activation of Ca^{2+} sensitive signal transduction pathways, such as calcineurin pathway. Calcineurin is a Ca^{2+} /calmodulin-dependent phosphatase that regulates hypertrophic response (Molkentin *et al.*, 1998; Molkentin, 2000; Diedrichs *et al.*, 2004; Wilkins *et al.*, 2004). Once activated, calcineurin directly dephosphorylates members of nuclear factor of activated T-cells transcription factor family (NFATc3) in the cytoplasm, resulting in their nuclear translocation and activation of hypertrophic genes. In the nucleus, NFAT interacts specifically with a cardiac-restricted zinc finger protein GATA-4 involved in pathological cardiac hypertrophic response (Molkentin *et al.*, 1998). Calcineurin activity and expression is increased in failing hearts and it is paralleled by increased NFATc3 translocation to the nucleus and higher GATA-4 expression levels (Diedrichs *et al.*, 2004), with consequent reactivation of fetal genes (Molkentin *et al.*, 1998). Although pharmacological and genetic inhibition of calcineurin or NFAT in rodents suffice for regression of pathological hypertrophy (Sussman *et al.*, 1998; Meguro *et al.*, 1999; Lim *et al.*, 2000; Bueno *et al.*, 2002; Wilkins *et*

al., 2002; Bourajjaj *et al.*, 2008), to date it is unknown whether exercise training, a physiological stimulus, is able to deactivate calcineurin pathway associated with an anti-remodelling effect in heart failure.

In contrast to the proposed pathological role for calcineurin/NFAT signalling in the heart, insulin-like growth factor (IGF-I)/phosphatidylinositol-3 kinase (PI3K)/protein kinase B (Akt)/mammalian target of rapamycin (mTOR) pathway has been reported to mediate physiological hypertrophy associated with exercise training while it is deactivated in pressure-overload hypertrophy (Kemi *et al.*, 2008b).

Here, we investigated the effect of moderate-intensity exercise training on calcineurin and AKT/mTOR signalling pathways in a genetic model of sympathetic hyperactivity-induced heart failure (α_{2A}/α_{2C} ARKO). α_{2A}/α_{2C} ARKO mice display severe cardiac dysfunction associated hypertrophy, cardiac Ca^{2+} -handling abnormalities and clinical signs of heart failure by 7 months of age (Bartholomeu *et al.*, 2008; Ferreira *et al.*, 2008). Previously, we have demonstrated the beneficial effects of exercise training on cardiac function, Ca^{2+} -handling, and survival in α_{2A}/α_{2C} ARKO mice (Rolim *et al.*, 2007). Therefore, the hypotheses of the present study were that moderated exercise training in α_{2A}/α_{2C} ARKO mice would: 1) deactivate the calcineurin signal pathway, 2) activate Akt/mTOR signal pathway and, 3) reduce cardiac mass and cardiac myocyte dimensions associated with a reduced expression of fetal genes.

Methods

Sampling. A cohort of male congenic α_{2A}/α_{2C} ARKO mice in a C57BL6/J genetic background and their wild-type controls (WT) were studied from 5 to 7 months of age. At 7 months of age, α_{2A}/α_{2C} ARKO mice present severe cardiac dysfunction associated with exercise intolerance and increased mortality rate (Brum *et al.*, 2002; Rolim *et al.*, 2007; Bartholomeu *et al.*, 2008). Mice were maintained in a 12:12h light-dark cycle and temperature-controlled environment (22°C) with free access to standard laboratory chow (Nuvital Nutrientes, Curitiba, PR Brazil) and tap water. This study was conducted in accordance with the ethical principles in animal research adopted by the Brazilian College of Animal Experimentation (www.cobea.org.br). The animal care and protocols in this

study were reviewed and approved by the Ethical Committee of Medical School of University of Sao Paulo (174/06).

Graded treadmill exercise test. Exercise capacity, estimated by total distance run, was evaluated with a graded treadmill exercise protocol for mice as previously described (Ferreira *et al.*, 2007). Briefly, after being adapted to treadmill exercises over 1 wk (10 min each session), mice were placed in the exercise streak and allowed to acclimatize for at least 30 min. Exercise intensity was increased by 3m/min (6-33m/min) every 3 min at 0% grade until exhaustion. The graded treadmill exercise test was performed in WT and α_{2A}/α_{2C} ARKO before and after experimental protocol to estimate exercise capacity, and at the fourth week of training in α_{2A}/α_{2C} ARKO to readjust exercise training intensity.

Exercise training protocol. α_{2A}/α_{2C} ARKO mice performed moderated intensity exercise training on a motor treadmill over 8 wk (from 5th-7th month of age), 5 days/wk. The running speed and duration of exercise were progressively increased to elicit 60% of maximal speed at the second week of training. At the fourth week of training, run capacity was evaluated in order to readjust exercise training intensity. In untrained α_{2A}/α_{2C} ARKO and WT mice, treadmill running skills were maintained by treadmill running for 5 min, three times a week. This procedure was also performed in order to avoid any interference of treadmill stress on the variables studied. This latter activity did not seem to alter maximal exercise capacity (Fig. 1A).

Cardiovascular measurements. Noninvasive cardiac function was assessed by two-dimensional guided M-mode echocardiography, in halothane-anesthetized WT and α_{2A}/α_{2C} ARKO mice before and after experimental protocol. Briefly, mice were positioned in the supine position with front paws wide open, and an ultrasound transmission gel was applied to the precordium. Transthoracic echocardiography was performed using an Acuson Sequoia model 512 echocardiographer equipped with a 14-MHz linear transducer. The left ventricular end-diastolic and end-systolic dimension (LVEDD and LVESD,

respectively), and the left ventricle systolic function by fractional shortening (FS) were evaluated. The FS was estimated as follows:

$$\text{FS\%} = \frac{\text{LVEDD} - \text{LVESD}}{\text{LVEDD}} \times 100$$

Structural analysis. Twenty-four hours after the last exercise training session, a subset of mice was killed by intravenous injection of sodium pentobarbital (120 mg/kg) and their tissues harvested. The heart was stopped at diastole (KCl, 14 mM) and dissected to obtain the left ventricle, which corresponds to the remaining organ upon removal of both atria and free wall of the right ventricle. For morphometric analysis, left ventricle samples obtained from the free wall, at the level of papillary muscle, were fixed in 4% buffered formalin and embedded in paraffin, cut in 4 μm sections and subsequently stained with hematoxylin and eosin. Two randomly selected sections from each animal were visualized by light microscopy using an objective with a calibrated magnification (400x). Myocytes with visible nucleus and intact cellular membranes were chosen for diameter determination. The width of individually isolated cardiomyocyte displayed on a viewing screen was manually traced, across the middle of the nuclei, with a digitizing pad and determined by a computer assisted image analysis system (Quantimet 520; Cambridge Instruments, UK). For each animal approximately 15 visual fields were analyzed.

RT-PCR. α_{2A}/α_{2C} ARKO mice present reactivation of fetal genes involved in cardiac remodelling and failure (Bartholomeu *et al.*, 2008). Therefore, we further evaluated whether exercise training would decrease β -myosin heavy chain (β -MHC) gene expression. RNA was isolated from left ventricle tissue with Trizol reagent (GIBCO-BRL, Invitrogen Corp., Carlsbad, CA, USA). cDNA was synthesized using Superscript III Reverse Transcriptase (200 U/ml, Invitrogen) at 42°C for 50 min. Taq DNA Polymerase (Fermentas, Burlington, ON, Canada) was used for DNA amplification in the presence of specific primers for cardiac isoform of β -MHC. The specific primer sequences were: β -MHC sense, 5' TGGCAAGACGGTGACTGTG 3'; β -MHC antisense, 5' CTCAAGGAGCGCTACGCTT 3'. For each cDNA, the number of amplification cycles was that necessary for 50% saturation, as determined in preliminary assays. mRNA levels

were normalized to that of β -actin mRNA (which is not changed in this model of HF) in the same assay.

Cellular Fractionation. Immediately following the experimental protocol, mice were killed by cervical dislocation and hearts were minced and homogenized in ice-cold homogenization RIPA buffer (150mM NaCl, 0.5% deoxycholate, 1% Triton100, 1:300 Sigma protease inhibitor cocktail and 50mM Tris-HCl at pH 7.4). The homogenate was centrifuged at 100 x g for 5 min (4°C). The resulting supernatant was centrifuged again at 600 x g for 5 min (4°C) to obtain the nuclear pellet and the cytoplasmic extract (supernatant). The pellet was incubated with RIPA buffer containing 0.3% SDS and DNase (1mg/ml) for 30 min (4°C) and then centrifuged at 2000 x g for 10 min (4°C) to obtain the nuclear extract (supernatant).

Western blot analysis. Calcineurin B, NFATc3 and GATA-4 expression levels were evaluated by Western blotting in total, cytoplasmic and nuclear extracts from WT and α_{2A}/α_{2C} ARKO hearts. In addition, Akt, p-Akt ser473, mTOR , p-mTOR ser2448, Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII), p-CaMKII thr286, class II histone deacetylases HDAC4 and HDAC5 expression levels were evaluated in cytoplasmic extracts. Briefly, samples were subjected to SDS-PAGE in polyacrylamide gels (8-12%) depending on protein molecular weight. After electrophoresis, proteins were electrotransferred to nitrocellulose membrane (BioRad Biosciences; Piscataway, NJ, USA). Equal loading of samples and transfer efficiency were monitored with the use of 0.5% Ponceau S staining of the blot membrane. The blotted membrane was then blocked (5% nonfat dry milk, 10 mM Tris-HCl (pH = 7.6), 150 mM NaCl, and 0.1% Tween 20) for 2h at room temperature and then incubated overnight at 4°C with specific antibodies against calcineurin B (Upstate, NY, USA), NFATc3, GATA-4, Lamin B, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and troponin I (Santa Cruz Bio Inc., CA, USA), Akt, p-Akt Ser473, mTOR, p-mTOR ser2448, CaMKII, p-CaMKII thr286, HDAC4 and HDAC5 (Cell Signaling Tech., MA, USA). Binding of the primary antibody was detected with the use of peroxidase-conjugated secondary antibodies (rabbit or mouse, depending on the

protein, for 2h at room temperature) and developed using enhanced chemiluminescence (Amersham Biosciences, NJ, USA) detected by autoradiography. Quantification analysis of blots was performed with the use of Scion Image software (Scion based on NIH image). Samples were normalized to relative changes in Lamin B (nucleus) and GAPDH (cytoplasm) and expressed as percent of control.

Confocal Microscopy. Isolated cardiomyocytes were analyzed by confocal microscopy to compare the cellular localization of NFATc3 and GATA-4. Cardiomyocytes from WT and α_{2A}/α_{2C} ARKO mice were enzymatically isolated as previously described (Guatimosim *et al.*, 2001). Briefly, the hearts were mounted on a Langendorff system, perfused for ~5 min with calcium-free solution containing (in mM): 130 NaCl, 5.4 KCl, 0.5 MgCl₂, 0.33 NaH₂PO₄, 1 lactate, 3 pyruvate, 22 glucose and 25 HEPES (pH=7.4). Afterwards, the hearts were perfused for 10–15 min with a solution containing 1 mg/ml collagenase type II (Worthington, USA) and 0.17 mg/ml protease type IX (Sigma Chemicals Co., St. Louis, MO, USA). The digested heart was then removed from the cannula, and the ventricles were cut into small pieces. Single cells were isolated by mechanical trituration and stored in Dulbecco's modified Eagle's medium (Sigma Chemicals Co., St Louis, MO, USA) supplemented with 10% fetal bovine serum (Cultilab, Brazil). Only Ca²⁺tolerant, quiescent, rodshaped myocytes showing clear cross striations were studied. Cells were then fixed with 4% paraformaldehyde and permeabilized as previously described (Guatimosim *et al.*, 2008). Cardiomyocytes were stained with primary antibodies against NFATc3 and GATA-4 (1:100; Santa Cruz). Cells were then stained with secondary antibodies conjugated to Alexa (1:1000; Invitrogen), followed by visualization of signals by confocal imaging using the ZEISS Meta confocal microscope (Zeiss Germany) from CEMEL (Biological Sciences Institute, UFMG, Brazil).

Cyclosporine treatment. To test whether calcineurin signalling pathway would be functionally involved in ventricular dysfunction and remodelling in this heart failure animal model, α_{2A}/α_{2C} ARKO mice (6 months of age) were treated for 4 weeks with cyclosporine (Sandimmun, Novartis, NJ, USA) delivered once a day at a rate of 25mg/kg/day (i.p.). This

dose was previously reported to inhibit calcineurin phosphatase activity in mice (Meguro *et al.*, 1999).

Statistical analysis. Data are presented as means \pm SE. Two-way analysis of variance (ANOVA) with a *post-hoc* testing by Duncan (Statistica software, StatSoft, Inc., Tulsa, OK, USA) was used to compare the effect of genotype (WT and α_{2A}/α_{2C} ARKO mice) and exercise training (untrained and trained) on data from Figure 1. One-way analysis of variance (ANOVA) with a *post-hoc* testing by Duncan (Statistica software, StarSoft, Inc, Tulsa, OK, USA) was used to analyze data from Figures 2,3,4 and 5. Statistical significance was considered achieved when the value of P was < 0.05 .

Results

Exercise training in α_{2A}/α_{2C} ARKO improves cardiac function and exercise tolerance associated with a cardiac anti-remodelling effect

α_{2A}/α_{2C} -ARKO mice displayed exercise intolerance and lower basal FS when compared to WT control mice (Figure 1A, B). Exercise training increased cardiac function and exercise capacity in α_{2A}/α_{2C} ARKO mice to WT levels. As expected, exercise training increased exercise capacity of WT mice with no changes in FS, heart weight and cardiomyocyte width. Consistent with left ventricle dysfunction, α_{2A}/α_{2C} -ARKO mice presented increased heart weight, cardiomyocyte width, and left ventricular dilation linked to increased LVESD and LVEDD (Fig. 1C and Table 1). Eight weeks of exercise training induced a significant cardiac anti-remodelling effect, reducing heart weight and cardiomyocyte width of α_{2A}/α_{2C} ARKO (Figure 1). In addition, exercise training prevented lung water retention observed in α_{2A}/α_{2C} ARKO mice (Table 1).

Exercise training reduces calcineurin-mediated signalling pathway activation in α_{2A}/α_{2C} ARKO failing hearts

Since calcineurin activation plays an important role in pathological cardiac hypertrophy (Molkentin *et al.*, 1998; Molkentin, 2000; Diedrichs *et al.*, 2004; Wilkins *et al.*, 2004), we evaluated expression levels of calcineurin B and its downstream targets, NFATc3 and GATA-4 in hearts of WT and α_{2A}/α_{2C} ARKO mice. No changes in cardiac calcineurin B total levels were observed in untrained and trained α_{2A}/α_{2C} ARKO compared with WT mice (Figure 2A, B). Nuclear calcineurin B translocation was significantly increased in untrained α_{2A}/α_{2C} ARKO, whereas exercise training decreased it toward WT levels (Figure 2A, B).

Even though total cardiac NFATc3 expression levels were similar among all three groups studied (Figure 2A,C), we observed a significant increase of NFATc3 translocation to nucleus in α_{2A}/α_{2C} ARKO (Figure 2A,C). Interestingly, exercise training decreased nuclear expression of NFATc3 to WT levels, which suggests an inhibition of NFATc3 nuclear translocation after exercise training in α_{2A}/α_{2C} ARKO. Total and nuclear GATA-4 expression levels were elevated in α_{2A}/α_{2C} ARKO compared to WT mice (Figure 2A,D). Exercise training reduced both GATA-4 expression levels and nuclear translocation in α_{2A}/α_{2C} ARKO mice to WT levels. Immunofluorescence experiments corroborated these findings in isolated ventricular myocytes (Figure 2E).

The increased cardiac expression of GATA-4 and translocation of NFATc3 to nucleus was paralleled by increased β -MHC mRNA levels (13%, $p < 0.05$) in α_{2A}/α_{2C} ARKO compared with WT mice. Of interest, exercise trained α_{2A}/α_{2C} ARKO significantly reduced cardiac β -MHC gene expression (by 95%, $p < 0.05$) to levels similar to WT mice.

While calcineurin is a high-affinity target of Ca^{2+} /calmodulin, CaMKII is a low-affinity target of Ca^{2+} /calmodulin (Saucerman & Bers, 2008; Song *et al.*, 2008) also involved in cardiac hypertrophy and dilation (Maier, 2009). Therefore, we also evaluated the relative contribution of CaMKII signalling pathway in cardiac remodelling of α_{2A}/α_{2C} ARKO mice. No changes in CaMKII pathway activation were observed in α_{2A}/α_{2C} ARKO mice. CaMKII and p-CaMKII expression levels as well as HDAC4 and HDAC5 cytoplasmic levels were similar among all mice studied (Fig 3A-C). These results

suggest that CaMKII signalling pathway is not involved in exercise training anti-cardiac remodelling effect.

To further investigate whether a pro-survival Akt/mTOR pathway was involved in the cardiac anti-remodelling effect of exercise training in α_{2A}/α_{2C} ARKO mice, we evaluated expression levels of Akt and mTOR in heart homogenates from WT and α_{2A}/α_{2C} ARKO mice. Both cardiac Akt and mTOR expression and their phosphorylation levels at serine 473 and serine 2448, respectively, were similar among all mice studied (Figure 3D, E). These results indicate that beneficial effects exerted by exercise training are not related to Akt/mTOR signalling pathway activation.

Sustained calcineurin B inhibition reestablishes cardiac function and remodelling in α_{2A}/α_{2C} ARKO

To evaluate the relative contribution of calcineurin signalling pathway on ventricular dysfunction and remodelling in this heart failure model, we treated α_{2A}/α_{2C} ARKO mice with cyclosporine. As depicted in Figure 4, cyclosporine treatment improved FS (Figure 4A) and exercise capacity (Figure 4B) of α_{2A}/α_{2C} ARKO mice to WT values. The increased left ventricular function was associated with a pronounced cardiac anti-remodelling effect, characterized by reduced heart weight (Figure 4C) and LVEDD (Table 1). As expected, 4 week-cyclosporine treatment diminished nuclear translocation of calcineurin B, NFATc3 and GATA-4 in α_{2A}/α_{2C} ARKO mice to WT levels (Figure 5 A-D). No changes in pro-survival Akt/mTOR pathway were observed in cyclosporine treated α_{2A}/α_{2C} ARKO mice (Figure 5 E, F).

Discussion

Exercise training is a key intervention for prevention and therapy in cardiology (Roveda *et al.*, 2003; Jonsdottir *et al.*, 2006; Wisloff *et al.*, 2007), and its effect on cardiac function and structure in heart failure includes improved net balance of Ca²⁺-handling proteins, improved left ventricular function and cardiac anti-remodelling effect (Lu *et al.*,

2002; Rolim *et al.*, 2007; Kemi *et al.*, 2008a; Medeiros *et al.*, 2008). However, the mechanisms underlying exercise training-induced cardiac anti-remodelling effect in heart failure remains unknown. The key findings of the present study are that moderate-intensity exercise training reduced nuclear translocation of calcineurin B, NFATc3 and GATA-4 without significant changes of CaMKII signalling pathway in hearts of α_{2A}/α_{2C} ARKO mice. These molecular changes were paralleled by diminished cardiac mass and reduced expression of fetal cardiac genes. Indeed, exercise training in α_{2A}/α_{2C} ARKO mice improved exercise tolerance and ventricular function while reduced lung water retention. Unexpectedly, these changes occurred without concomitant activation of Akt-mTOR signalling pathway in this genetic model of heart failure.

Considering that both calcineurin and CaMKII are targets of Ca^{2+} /calmodulin, one might expect that both pathways could be involved in hypertrophic response of cardiac myocytes in α_{2A}/α_{2C} ARKO mice. However, the preferential activation of calcineurin pathway in our heart failure model might be explained by the higher affinity of Ca^{2+} -calmodulin for calcineurin than CaMKII. In fact, Song *et al.* (2008) demonstrated that differential Ca^{2+} -calmodulin binding affinities of targets (e.g calcineurin vs. CaMKII) predict the selective activation of distinct Ca^{2+} signaling pathways in paced adult rabbit ventricular myocytes (Song *et al.*, 2008).

Cardiac hypertrophy in α_{2A}/α_{2C} ARKO mice can be counteracted by cyclosporine treatment, which reinforces the involvement of calcineurin pathway on cardiac remodelling in this heart failure model. As the therapeutic use of cyclosporine for reducing heart failure-associated cardiac remodelling has been tempered by its known side effects (Molkentin, 2000), moderate intensity aerobic exercise training emerges as an adjuvant therapy for cardiac anti-remodelling effect by decreasing calcineurin/NFAT signalling pathway that is exacerbated in heart failure. In fact, Konhilas *et al.* (2006) reported that voluntary exercise decreased NFAT activity and reversed cardiac disease phenotype in hypertrophic cardiomyopathy animal model (Konhilas *et al.*, 2006). Our study extends the knowledge that exercise training deactivates calcineurin/NFAT signalling pathway to heart failure. This is particularly important since heart failure is a common endpoint of several cardiomyopathies.

The mechanisms by which exercise training decreases calcineurin/NFAT signalling pathways in heart failure might involve improved cardiac net balance of Ca^{2+} -handling proteins. In fact, we previously reported that moderate intensity aerobic exercise training improved cardiac intracellular Ca^{2+} regulation of α_{2A}/α_{2C} ARKO mice in different stages of heart failure by increasing the expression of sarcoplasmic reticulum Ca^{2+} -ATPase (SERCA2), and phosphorylation of phospholamban at both Serine16 and Threonine17 residues (Rolim *et al.*, 2007; Medeiros *et al.*, 2008). The impact of exercise training on SERCA2 and phosphorylation of phospholamban is positive for heart failure since both are decreased in human heart failure and associated with increased calcineurin phosphatase activity (Munch *et al.*, 2002). An alternative mechanism associated with decreased calcineurin signalling pathway by exercise training might be related to its effect on sympathetic nerve activity and neurohormones. Sympathetic hyperactivity, presently achieved by the use of genetically engineered mice, plays a prominent role in cardiac remodelling and failure (Barki-Harrington *et al.*, 2004) and β -adrenoceptor stimulation increases calcineurin activity in hearts of hypertensive rats (MacDonnell *et al.*, 2007). Exercise training is efficient in reducing sympathetic nerve activity of heart failure patients (Roveda *et al.*, 2003; Fraga *et al.*, 2007). In addition, we have also observed decreased circulating noradrenaline (Medeiros *et al.*, 2008), and cardiac angiotensin II (Pereira *et al.*, 2009) levels in exercise trained α_{2A}/α_{2C} ARKO mice.

Exercise training-induced deactivation of calcineurin signalling pathway in α_{2A}/α_{2C} ARKO mice was associated with improved ventricular function, reduced reactivation of cardiac fetal genes and decreased cardiomyocyte width, which highlights the role of exercise training in reversing pathological hypertrophy associated with heart failure in our model. However, one could expect that exercise training instead of deactivating a pathological pathway in cardiac remodelling would rather activate intracellular pathways involved in physiological cardiac hypertrophy, such as Akt/mTOR signalling pathway. Akt/mTOR signalling pathway has been reported as a key mediator of cardiac physiological growth, since constitutive activation of Akt leads to physiological hypertrophy and increased cardiac contractility (Condorelli *et al.*, 2002; McMullen *et al.*, 2003). Moreover,

physiological cardiac hypertrophy is blunted in Akt-1 knockout mice (DeBosch *et al.*, 2006).

We observed no changes in Akt/mTOR expression levels, as well as, in their phosphorylated levels in α_{2A}/α_{2C} ARKO when compared with WT and exercise trained α_{2A}/α_{2C} ARKO mice. This result was somehow unexpected but factors, such as exercise training intensity may contribute to this response. In fact, in parallel to observed skeletal muscle hypertrophy associated with Akt/mTOR signalling activation in high- but not low-intensity exercise training (Nader & Esser, 2001; Atherton *et al.*, 2005), exercise-induced physiological cardiac hypertrophy has been primarily observed in high-intensity exercise training regimens (McMullen *et al.*, 2003; Wilkins *et al.*, 2004; McMullen *et al.*, 2007; Kemi *et al.*, 2008a). Another factor worth mentioning is that the effect of exercise training on cardiac Akt/mTOR signaling pathway has been mainly studied in hearts of animals with preserved function (Kemi *et al.*, 2008a), but not in animal models of cardiovascular disease. Therefore, we cannot exclude that Akt/mTOR signalling pathway is not the main mechanism involved in cardiac anti-remodelling effect of exercise training in failing hearts.

In the present study, we took advantage of a genetic model of sympathetic hyperactivity-induced HF to assess the effect of moderate intensity aerobic exercise training on cardiac structure and function and the involvement of calcineurin/NFAT and Akt/mTOR signalling pathways in these responses. We cannot exclude that some of the results reported here may be influenced by exercise training intensity and stage of cardiomyopathy in α_{2A}/α_{2C} ARKO mice, but within this time window, moderate intensity aerobic exercise training in severe heart failure was effective in improving cardiac function with a cardiac anti-remodelling effect associated with deactivation of calcineurin/NFAT signalling pathway. It will be important, however, to further explore whether more intense exercise training regimen and earlier heart failure stages lead to different efficacy in reducing or preventing cardiac remodelling.

Conclusion

In summary, moderate intensity aerobic exercise training induced a cardiac anti-remodelling effect in heart failure mice associated with deactivation of calcineurin/NFAT

signalling pathway and decreased reactivation of fetal genes. These findings support the notion that deactivation of pathological hypertrophy signalling pathways is as preferential mechanism underlying cardiac anti-remodelling effect of moderate intensity aerobic exercise training in heart failure.

Acknowledgments

The authors want to thank the Fundação de Amparo à Pesquisa do Estado de São Paulo, São Paulo - SP (FAPESP # 2006/61523-7) for funding this present investigation. R.S.F.O. held a master's fellowship from FAPESP (# 2006/054826-3). P.C.B. holds scholarship from Conselho Nacional de Pesquisa e Desenvolvimento – Brasil (CNPq, BPQ). S.G. holds scholarship from Instituto do Milenio Toxins/MCT, FAPEMIG and CNPq.

References

- Atherton PJ, Babraj J, Smith K, Singh J, Rennie MJ & Wackerhage H. (2005). Selective activation of AMPK-PGC-1alpha or PKB-TSC2-mTOR signaling can explain specific adaptive responses to endurance or resistance training-like electrical muscle stimulation. *FASEB J* **19**, 786-788.
- Balke CW & Shorofsky SR. (1998). Alterations in calcium handling in cardiac hypertrophy and heart failure. *Cardiovasc Res* **37**, 290-299.
- Barki-Harrington L, Perrino C & Rockman HA. (2004). Network integration of the adrenergic system in cardiac hypertrophy. *Cardiovasc Res* **63**, 391-402.
- Bartholomeu JB, Vanzelli AS, Rolim NP, Ferreira JC, Bechara LR, Tanaka LY, Rosa KT, Alves MM, Medeiros A, Mattos KC, Coelho MA, Irigoyen MC, Krieger EM, Krieger JE, Negrao CE, Ramires PR, Guatimosim S & Brum PC. (2008). Intracellular mechanisms of specific beta-adrenoceptor antagonists involved in improved cardiac function and survival in a genetic model of heart failure. *J Mol Cell Cardiol* **45**, 240-249.
- Bourajjaj M, Armand AS, da Costa Martins PA, Weijts B, van der Nagel R, Heeneman S, Wehrens XH & De Windt LJ. (2008). NFATc2 is a necessary mediator of calcineurin-dependent cardiac hypertrophy and heart failure. *The Journal of biological chemistry*.
- Brum PC, Kosek J, Patterson A, Bernstein D & Kobilka B. (2002). Abnormal cardiac function associated with sympathetic nervous system hyperactivity in mice. *Am J Physiol Heart Circ Physiol* **283**, H1838-1845.
- Bueno OF, Wilkins BJ, Tymitz KM, Glascock BJ, Kimball TF, Lorenz JN & Molkentin JD. (2002). Impaired cardiac hypertrophic response in Calcineurin Abeta -deficient mice. *Proc Natl Acad Sci U S A* **99**, 4586-4591.
- Bustamante JO, Ruknudin A & Sachs F. (1991). Stretch-activated channels in heart cells: relevance to cardiac hypertrophy. *J Cardiovasc Pharmacol* **17 Suppl 2**, S110-113.
- Condorelli G, Drusco A, Stassi G, Bellacosa A, Roncarati R, Iaccarino G, Russo MA, Gu Y, Dalton N, Chung C, Latronico MV, Napoli C, Sadoshima J, Croce CM & Ross J, Jr. (2002). Akt induces enhanced myocardial contractility and cell size in vivo in transgenic mice. *Proc Natl Acad Sci U S A* **99**, 12333-12338.
- DeBosch B, Treskov I, Lupu TS, Weinheimer C, Kovacs A, Courtois M & Muslin AJ. (2006). Akt1 is required for physiological cardiac growth. *Circulation* **113**, 2097-2104.

- Diedrichs H, Chi M, Boelck B, Mehlhorn U & Schwinger RH. (2004). Increased regulatory activity of the calcineurin/NFAT pathway in human heart failure. *Eur J Heart Fail* **6**, 3-9.
- Ferreira JC, Bacurau AV, Evangelista FS, Coelho MA, Oliveira EM, Casarini DE, Krieger JE & Brum PC. (2008). The role of local and systemic renin angiotensin system activation in a genetic model of sympathetic hyperactivity-induced heart failure in mice. *Am J Physiol Regul Integr Comp Physiol* **294**, R26-32.
- Ferreira JC, Rolim NP, Bartholomeu JB, Gobatto CA, Kokubun E & Brum PC. (2007). Maximal lactate steady state in running mice: effect of exercise training. *Clin Exp Pharmacol Physiol* **34**, 760-765.
- Fraga R, Franco FG, Roveda F, de Matos LN, Braga AM, Rondon MU, Rotta DR, Brum PC, Barretto AC, Middlekauff HR & Negrao CE. (2007). Exercise training reduces sympathetic nerve activity in heart failure patients treated with carvedilol. *Eur J Heart Fail* **9**, 630-636.
- Frey N & Olson EN. (2003). Cardiac hypertrophy: the good, the bad, and the ugly. *Annu Rev Physiol* **65**, 45-79.
- Guatimosim S, Amaya MJ, Guerra MT, Aguiar CJ, Goes AM, Gomez-Viquez NL, Rodrigues MA, Gomes DA, Martins-Cruz J, Lederer WJ & Leite MF. (2008). Nuclear Ca²⁺ regulates cardiomyocyte function. *Cell Calcium* **44**, 230-242.
- Guatimosim S, Sobie EA, dos Santos Cruz J, Martin LA & Lederer WJ. (2001). Molecular identification of a TTX-sensitive Ca(2+) current. *Am J Physiol Cell Physiol* **280**, C1327-1339.
- Gwathmey JK, Copelas L, MacKinnon R, Schoen FJ, Feldman MD, Grossman W & Morgan JP. (1987). Abnormal intracellular calcium handling in myocardium from patients with end-stage heart failure. *Circulation research* **61**, 70-76.
- Ho KK, Pinsky JL, Kannel WB & Levy D. (1993). The epidemiology of heart failure: the Framingham Study. *J Am Coll Cardiol* **22**, 6A-13A.
- Jonsdottir S, Andersen KK, Sigurosson AF & Sigurosson SB. (2006). The effect of physical training in chronic heart failure. *Eur J Heart Fail* **8**, 97-101.
- Kemi OJ, Ceci M, Condorelli G, Smith GL & Wisloff U. (2008a). Myocardial sarcoplasmic reticulum Ca²⁺ ATPase function is increased by aerobic interval training. *Eur J Cardiovasc Prev Rehabil* **15**, 145-148.

- Kemi OJ, Ceci M, Wisloff U, Grimaldi S, Gallo P, Smith GL, Condorelli G & Ellingsen O. (2008b). Activation or inactivation of cardiac Akt/mTOR signaling diverges physiological from pathological hypertrophy. *J Cell Physiol* **214**, 316-321.
- Konhilas JP, Watson PA, Maass A, Boucek DM, Horn T, Stauffer BL, Luckey SW, Rosenberg P & Leinwand LA. (2006). Exercise can prevent and reverse the severity of hypertrophic cardiomyopathy. *Circulation research* **98**, 540-548.
- Levy D, Garrison RJ, Savage DD, Kannel WB & Castelli WP. (1990). Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med* **322**, 1561-1566.
- Lim HW, De Windt LJ, Steinberg L, Taigen T, Witt SA, Kimball TR & Molkenin JD. (2000). Calcineurin expression, activation, and function in cardiac pressure-overload hypertrophy. *Circulation* **101**, 2431-2437.
- Lorell BH & Carabello BA. (2000). Left ventricular hypertrophy: pathogenesis, detection, and prognosis. *Circulation* **102**, 470-479.
- Lu L, Mei DF, Gu AG, Wang S, Lentzner B, Gutstein DE, Zwas D, Homma S, Yi GH & Wang J. (2002). Exercise training normalizes altered calcium-handling proteins during development of heart failure. *J Appl Physiol* **92**, 1524-1530.
- MacDonnell SM, Kubo H, Harris DM, Chen X, Berretta R, Barbe MF, Kolwicz S, Reger PO, Eckhart A, Renna BF, Koch WJ, Houser SR & Libonati JR. (2007). Calcineurin inhibition normalizes beta-adrenergic responsiveness in the spontaneously hypertensive rat. *Am J Physiol Heart Circ Physiol* **293**, H3122-3129.
- Maier LS. (2009). Role of CaMKII for signaling and regulation in the heart. *Front Biosci* **14**, 486-496.
- McMullen JR, Amirahmadi F, Woodcock EA, Schinke-Braun M, Bouwman RD, Hewitt KA, Mollica JP, Zhang L, Zhang Y, Shioi T, Buerger A, Izumo S, Jay PY & Jennings GL. (2007). Protective effects of exercise and phosphoinositide 3-kinase(p110alpha) signaling in dilated and hypertrophic cardiomyopathy. *Proc Natl Acad Sci U S A* **104**, 612-617.
- McMullen JR, Shioi T, Zhang L, Tarnavski O, Sherwood MC, Kang PM & Izumo S. (2003). Phosphoinositide 3-kinase(p110alpha) plays a critical role for the induction of physiological, but not pathological, cardiac hypertrophy. *Proc Natl Acad Sci U S A* **100**, 12355-12360.
- Medeiros A, Rolim NP, Oliveira RS, Rosa KT, Mattos KC, Casarini DE, Irigoyen MC, Krieger EM, Krieger JE, Negrao CE & Brum PC. (2008). Exercise training delays

- cardiac dysfunction and prevents calcium handling abnormalities in sympathetic hyperactivity-induced heart failure mice. *J Appl Physiol* **104**, 103-109.
- Meguro T, Hong C, Asai K, Takagi G, McKinsey TA, Olson EN & Vatner SF. (1999). Cyclosporine attenuates pressure-overload hypertrophy in mice while enhancing susceptibility to decompensation and heart failure. *Circulation research* **84**, 735-740.
- Molkentin JD. (2000). Calcineurin and beyond: cardiac hypertrophic signaling. *Circulation research* **87**, 731-738.
- Molkentin JD, Lu JR, Antos CL, Markham B, Richardson J, Robbins J, Grant SR & Olson EN. (1998). A calcineurin-dependent transcriptional pathway for cardiac hypertrophy. *Cell* **93**, 215-228.
- Munch G, Bolck B, Karczewski P & Schwinger RH. (2002). Evidence for calcineurin-mediated regulation of SERCA 2a activity in human myocardium. *J Mol Cell Cardiol* **34**, 321-334.
- Nader GA & Esser KA. (2001). Intracellular signaling specificity in skeletal muscle in response to different modes of exercise. *J Appl Physiol* **90**, 1936-1942.
- Naylor LH, George K, O'Driscoll G & Green DJ. (2008). The athlete's heart: a contemporary appraisal of the 'Morganroth hypothesis'. *Sports Med* **38**, 69-90.
- Pereira MG, Ferreira JC, Bueno CR, Jr., Mattos KC, Rosa KT, Irigoyen MC, Oliveira EM, Krieger JE & Brum PC. (2009). Exercise training reduces cardiac angiotensin II levels and prevents cardiac dysfunction in a genetic model of sympathetic hyperactivity-induced heart failure in mice. *Eur J Appl Physiol*.
- Rolim NP, Medeiros A, Rosa KT, Mattos KC, Irigoyen MC, Krieger EM, Krieger JE, Negrao CE & Brum PC. (2007). Exercise training improves the net balance of cardiac Ca²⁺ handling protein expression in heart failure. *Physiol Genomics* **29**, 246-252.
- Rossmann EI, Petre RE, Chaudhary KW, Piacentino V, 3rd, Janssen PM, Gaughan JP, Houser SR & Margulies KB. (2004). Abnormal frequency-dependent responses represent the pathophysiologic signature of contractile failure in human myocardium. *J Mol Cell Cardiol* **36**, 33-42.
- Roveda F, Middlekauff HR, Rondon MU, Reis SF, Souza M, Nastari L, Barretto AC, Krieger EM & Negrao CE. (2003). The effects of exercise training on sympathetic neural activation in advanced heart failure: a randomized controlled trial. *J Am Coll Cardiol* **42**, 854-860.

- Saucerman JJ & Bers DM. (2008). Calmodulin mediates differential sensitivity of CaMKII and calcineurin to local Ca²⁺ in cardiac myocytes. *Biophysical journal* **95**, 4597-4612.
- Schwinger RH, Bohm M, Schmidt U, Karczewski P, Bavendiek U, Flesch M, Krause EG & Erdmann E. (1995). Unchanged protein levels of SERCA II and phospholamban but reduced Ca²⁺ uptake and Ca(2+)-ATPase activity of cardiac sarcoplasmic reticulum from dilated cardiomyopathy patients compared with patients with nonfailing hearts. *Circulation* **92**, 3220-3228.
- Seki S, Nagai M, Takeda H, Onodera T, Okazaki F, Taniguchi M, Taniguchi I & Mochizuki S. (2003). Impaired Ca²⁺ handling in perfused hypertrophic hearts from Dahl salt-sensitive rats. *Hypertens Res* **26**, 643-653.
- Song Q, Saucerman JJ, Bossuyt J & Bers DM. (2008). Differential integration of Ca²⁺-calmodulin signal in intact ventricular myocytes at low and high affinity Ca²⁺-calmodulin targets. *The Journal of biological chemistry* **283**, 31531-31540.
- Sussman MA, Lim HW, Gude N, Taigen T, Olson EN, Robbins J, Colbert MC, Gualberto A, Wiecek DF & Molkentin JD. (1998). Prevention of cardiac hypertrophy in mice by calcineurin inhibition. *Science* **281**, 1690-1693.
- Wilkins BJ, Dai YS, Bueno OF, Parsons SA, Xu J, Plank DM, Jones F, Kimball TR & Molkentin JD. (2004). Calcineurin/NFAT coupling participates in pathological, but not physiological, cardiac hypertrophy. *Circulation research* **94**, 110-118.
- Wilkins BJ, De Windt LJ, Bueno OF, Braz JC, Glascock BJ, Kimball TF & Molkentin JD. (2002). Targeted disruption of NFATc3, but not NFATc4, reveals an intrinsic defect in calcineurin-mediated cardiac hypertrophic growth. *Mol Cell Biol* **22**, 7603-7613.
- Wisloff U, Stoylen A, Loennechen JP, Bruvold M, Rognum O, Haram PM, Tjonna AE, Helgerud J, Slordahl SA, Lee SJ, Videm V, Bye A, Smith GL, Najjar SM, Ellingsen O & Skjaerpe T. (2007). Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation* **115**, 3086-3094.

Author contributions

R.S.F.O. and J.C.B.F. contributed to study design and performed most part of experiments. E.R.M.G. and S.G. collaborated on experiments shown in Figure 2E, N.A.P. collaborated on real time PCR experiments, A.M. and N.P.L.R collaborated on fractional shortening evaluation and western blotting experiments. P.C.B. directed and designed the study, and wrote the manuscript. S.G. revised the manuscript critically.

Table 1: Structural parameters in WT and α_{2A}/α_{2C} ARKO mice.

	WT	KO	KOt	KOc
LVEDD, mm	3.74±0.08 (8)	4.03±0.06 (10)*	3.98±0.03 (10)*	3.84±0.11 (9)
LVESD, mm	3.07±0.08 (8)	3.44±0.06 (10)*	3.22±0.04 (10)	3.08±0.10 (9)#
AT, mg/mm	0.42±0.03 (10)	0.43±0.04 (8)	0.39±0.03 (13)	0.34±0.02 (8)
RV, mg/mm	1.60±0.16 (10)	1.52±0.06 (8)	1.38±0.13 (13)	1.45±0.09 (8)
LV, mg/mm	5.64±0.22 (10)	6.52±0.25 (8)*	5.84±0.20 (13)	5.95±0.13 (8)
Lung wet/dry ratio	5.57±0.13 (13)	6.44±0.36 (7)*	5.27±0.19 (16)#	5.52±0.21 (9)#

LVEDD, left ventricular diastolic diameter; LVESD, left ventricular systolic diameter, AT, atria; RV, right ventricle; LV, left ventricle. AT, RV and LV weights were normalized by tibial length. Data are presented as mean ±SE.* $P < 0.05$ vs. WT; #, $P < 0.05$ vs. KO. The number of animals studied in each group is showed between parentheses. Data were analyzed by one-way ANOVA with *post-hoc* testing by Duncan.

Legends

Figure 1. Exercise training reestablishes cardiac function and leads to a cardiac anti-remodelling effect in α_{2A}/α_{2C} ARKO.

Total distance run (A), fractional shortening (B), heart weight and cardiomyocyte width (C) measurements were performed in untrained wild-type (WT), trained wild-type (WTt), untrained α_{2A}/α_{2C} ARKO (KO) and trained α_{2A}/α_{2C} ARKO (KOt), before (\square) and after (\blacksquare) exercise training protocol. Note that untrained KO presented exercise intolerance, cardiac dysfunction and cardiac hypertrophy. Exercise training also improved cardiac function and decreased cardiac mass and cardiomyocyte width in α_{2A}/α_{2C} ARKO mice. Data are presented as mean \pm SE. * $P < 0.05$ vs. WT, # $P < 0.05$ vs. KO. Figures A and B were analyzed by two-way ANOVA for repeated measurements with *post-hoc* testing by Duncan. Figure C was analyzed by two-way ANOVA with *post-hoc* testing by Duncan.

Figure 2. Cardiac anti-remodelling effect of exercise training in α_{2A}/α_{2C} ARKO is associated to decreased calcineurin signalling pathway activation.

Representative blots of Calcineurin B, NFATc3 and GATA-4 in total and nuclear extracts from untrained wild type (WT), untrained α_{2A}/α_{2C} ARKO (KO) and trained α_{2A}/α_{2C} ARKO (KOt) after exercise training protocol (A). Total and nuclear extracts were normalized by troponin I and laminin B, respectively. Calcineurin B (B), NFATc3 (C) and GATA-4 (D) expression levels and nuclear/cytoplasmic ratio from WT, KO and KOt. E. Confocal images showing immunofluorescence labelled cardiomyocytes stained with anti-NFATc3 or anti-GATA-4 antibodies. Cytoplasmic and nuclear localization of NFATc3 and GATA-4 in untrained WT, untrained KO and trained KOt. Exercise training prevented nuclear calcineurin B, NFATc3 and GATA-4 translocation in α_{2A}/α_{2C} ARKO mice. Red arrows indicate cell nucleus. Bar = 10 μ m. Data are presented as mean \pm SE. * $P < 0.05$ vs. WT, # $P < 0.05$ vs. KOt. Data were analyzed by one-way ANOVA with *post-hoc* testing by Duncan.

Figure 3. Exercise training has no effect on CaMKII and Akt signalling pathway activation in α_{2A}/α_{2C} ARKO.

Representative blots of CaMKII, p-CaMKII thr286, HDAC4, HDAC5, Akt, p-Akt ser473, mTOR, p-mTOR ser2448 in cytoplasmic fraction from untrained wild type (WT), untrained α_{2A}/α_{2C} ARKO (KO) and trained α_{2A}/α_{2C} ARKO (KOt) after exercise training protocol (A). CaMKII and p-CaMKII thr286 (B), HDAC4 and HDAC5 (C), Akt and p-Akt ser473 (D), mTOR and p-mTOR ser2448 (E) expression levels in cytoplasmic fraction from WT, KO and KOt (G). Note that cardiac CaMKII, Akt and mTOR expression levels and their phosphorylation at threonine 286, serine 473 and serine 2448, respectively, were similar among all mice studied. Data are presented as mean \pm SE. Data were analyzed by one-way ANOVA with *post-hoc* testing by Duncan.

Figure 4. Sustained calcineurin B inhibition reestablishes cardiac function and leads to a cardiac anti-remodelling effect in α_{2A}/α_{2C} ARKO.

Total distance run (A), fractional shortening (B) and heart weight (C) from wild-type (WT), α_{2A}/α_{2C} ARKO (KO) and cyclosporine-treated α_{2A}/α_{2C} ARKO (KOc). Note that 4 weeks of cyclosporine treatment improved cardiac function in 28% and decreased heart weight to WT levels. No changes in exercise tolerance were observed in KO mice treated with cyclosporine. Data are presented as mean \pm SE. * $P<0.05$ vs. WT, # $P<0.05$ vs. KO. Data were analyzed by one-way ANOVA with *post-hoc* testing by Duncan.

Figure 5. Sustained cyclosporine treatment decreases calcineurin/NFAT pathway in α_{2A}/α_{2C} ARKO

Representative blots of Calcineurin B, NFATc3, GATA-4, in total and nuclear extracts and Akt, p-Akt ser473, mTOR and p-mTOR ser2448 in cytoplasmic extract from untrained wild type (WT), untrained α_{2A}/α_{2C} ARKO (KO) and cyclosporine-treated α_{2A}/α_{2C} ARKO (KOc) (A). Calcineurin B (B), NFATc3 (C) and GATA-4 (D) expression levels and nuclear/cytoplasmic ratio from WT, KO and KOc. Cytoplasmic Akt and p-Akt ser473 (E), mTOR and p-mTOR ser2448 (F) expression levels from WT, KO and KOc. Cyclosporine

treatment prevented nuclear cyclosporine B, NFATc3 and GATA-4 translocation in α_{2A}/α_{2C} ARKO mice. * $P < 0.05$ vs. WT, # $P < 0.05$ vs. KOc. Data were analyzed by one-way ANOVA with *post-hoc* testing by Duncan.





Figure 1. Cellulose and Shattering






Figure 2. Colville et al.




Figure 4 olive ra ct oil.




Figure 6. CDKN1B, CDKN1C,